



Bright Futures Parent Handout

4 Month Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

FAMILY FUNCTIONING

How Your Family Is Doing

- Take time for yourself.
- Take time together with your partner.
- Spend time alone with your other children.
- Encourage your partner to help care for your baby.
- Choose a mature, trained, and responsible babysitter or caregiver.
- You can talk with us about your child care choices.
- Hold, cuddle, talk to, and sing to your baby each day.
- Massaging your infant may help your baby go to sleep more easily.
- Get help if you and your partner are in conflict. Let us know. We can help.

Feeding Your Baby

- Feed only breast milk or iron-fortified formula in the first 4–6 months.

If Breastfeeding

- If you are still breastfeeding, that's great!
- Plan for pumping and storage of breast milk.

If Formula Feeding

- Make sure to prepare, heat, and store the formula safely.
- Hold your baby so you can look at each other while feeding.
- Do not prop the bottle.
- Do not give your baby a bottle in the crib.

Solid Food

- You may begin to feed your baby solid food when your baby is ready.
- Some of the signs your baby is ready for solids
 - Opens mouth for the spoon.
 - Sits with support.
 - Good head and neck control.
 - Interest in foods you eat.
- Avoid foods that cause allergy—peanuts, tree nuts, fish, and shellfish.
- Avoid feeding your baby too much by following the baby's signs of fullness

NUTRITIONAL ADEQUACY AND GROWTH

NUTRITION

- Leaning back
- Turning away
- Ask us about programs like WIC that can help get food for you if you are breastfeeding and formula for your baby if you are formula feeding.

Safety

- Use a rear-facing car safety seat in the back seat in all vehicles.
- Always wear a seat belt and never drive after using alcohol or drugs.
- Keep small objects and plastic bags away from your baby.
- Keep a hand on your baby on any high surface from which she can fall and be hurt.
- Prevent burns by setting your water heater so the temperature at the faucet is 120°F or lower.
- Do not drink hot drinks when holding your baby.
- Never leave your baby alone in bathwater, even in a bath seat or ring.
- The kitchen is the most dangerous room. Don't let your baby crawl around there; use a playpen or high chair instead.
- Do not use a baby walker.

SAFETY

Your Changing Baby

- Keep routines for feeding, nap time, and bedtime.

Crib/Playpen

- Put your baby to sleep on her back.
- In a crib that meets current safety standards, with no drop-side rail and slats no more than 2³/₈ inches apart. Find more information on the Consumer Product Safety Commission Web site at www.cpsc.gov.
- If your crib has a drop-side rail, keep it up and locked at all times. Contact the crib company to see if there is a device to keep the drop-side rail from falling down.

INFANT DEVELOPMENT

INFANT DEVELOPMENT

Playtime

- Learn what things your baby likes and does not like.
- Encourage active play.
 - Offer mirrors, floor gyms, and colorful toys to hold.
 - Tummy time—put your baby on his tummy when awake and you can watch.
- Promote quiet play.
 - Hold and talk with your baby.
 - Read to your baby often.

Crying

- Give your baby a pacifier or his fingers or thumb to suck when crying.

Healthy Teeth

- Go to your own dentist twice yearly. It is important to keep your teeth healthy so that you don't pass bacteria that causes tooth decay on to your baby.
- Do not share spoons or cups with your baby or use your mouth to clean the baby's pacifier.
- Use a cold teething ring if your baby has sore gums with teething.

ORAL HEALTH

What to Expect at Your Baby's 6 Month Visit

We will talk about

- Introducing solid food
- Getting help with your baby
- Home and car safety
- Brushing your baby's teeth
- Reading to and teaching your baby

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org

The Perkins Pediatric Clinic, LLC
Michael W. Perkins, MD
614 North Pine Street
DeRidder, LA 70634

phone: (337) 462-6000
www.perkinspediatrics.com





American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics. Updated 10/11. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.

Birth to 6 Months



BIRTH TO 6 MONTHS


Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries — most of which could be prevented?

Often, injuries happen because parents are not aware of what their children can do. Children *learn fast*, and before you know it, your child will be *wiggling* off a bed or *reaching* for your cup of hot coffee.

Car Injuries

Car crashes are a great threat to your child's life and health. Most injuries and deaths from car crashes **can be prevented** by the use of car safety seats. Your child, besides being much safer in a car safety seat, will behave better, so you can pay attention to your driving. Make your newborn's first ride home from the hospital a safe one — in a car safety seat. Your infant should ride in the back seat in a rear-facing car seat.




Make certain that your baby's car safety seat is installed correctly. Read and follow the instructions that come with the car safety seat and the sections in the owners' manual of your car on using car safety seats correctly. Use the car safety seat EVERY time your child is in the car.

NEVER put an infant in the front seat of a car with a passenger air bag.

Falls

Babies *wiggle* and *move* and *push* against things with their feet soon after they are born. Even these very first movements can result in a fall. As your baby grows and is able to roll over, he or she may fall off of things unless protected. **Do not leave your baby alone** on changing tables, beds, sofas, or chairs. **Put your baby in a safe place** such as a crib or playpen when you cannot hold him.



Your baby may be able to crawl as early as 6 months. **Use gates on stairways and close doors** to keep your baby out of rooms where he or she might get hurt. **Install operable window guards** on all windows above the first floor.

Do not use a baby walker. Your baby may tip the walker over, fall out of it, or fall down stairs and seriously injure his head. Baby walkers let children get to places where they can pull heavy objects or hot food on themselves.

If your child has a serious fall or does not act normally after a fall, call your doctor.

(over)

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™




Burns

At 3 to 5 months, babies will wave their fists and grab at things. **NEVER carry your baby and hot liquids, such as coffee, or foods at the same time.** Your baby can get burned. You can't handle both! To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.

If your baby gets burned, immediately put the burned area in cold water. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth and call your doctor.

To protect your baby from house fires, be sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.




Choking and Suffocation

Babies explore their environment by putting anything and everything into their mouths. **NEVER leave small objects in your baby's reach, even for a moment.** NEVER feed your baby hard pieces of food such as chunks of raw carrots, apples, hot dogs, grapes, peanuts, and popcorn. Cut all the foods you feed your baby into thin pieces to prevent choking. **Be prepared if your baby starts to choke. Ask your doctor to recommend the steps you need to know. Learn how to save the life of a choking child.**

To prevent possible suffocation and reduce the risk of sudden infant death syndrome (SIDS), **your baby should always sleep on his or her back. Your baby should have his or her own crib or bassinet with no pillows, stuffed toys, bumpers, or loose bedding. NEVER put your baby on a water bed, bean bag, or anything that is soft enough to cover the face and block air to the nose and mouth.**

Plastic wrappers and bags form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your baby.




From Your Doctor

The Perkins Pediatric Clinic, LLC
Michael W. Perkins, MD
614 North Pine Street
DeRidder, LA 70634
(337) 462-6000
www.perkinspediatrics.com

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on the individual facts and circumstances.

Starting Solid Foods



Rice, oatmeal, or barley? What infant cereal or other food will be on the menu for your baby's first solid meal? And have you set a date?

At this point, you may have a plan or are confused because you have received too much advice from family and friends with different opinions. To help you prepare for your baby's transition to solid food, read on for more information from the American Academy of Pediatrics (AAP).

When can my baby begin solid foods?

The following are some guidelines from the AAP book *Nutrition: What Every Parent Needs to Know*. Remember that each child's readiness depends on his own rate of development.

- **Can he hold his head up?** Your baby should be able to sit in a high chair, feeding seat, or infant seat with good head control.
- **Does he open his mouth when food comes his way?** Babies may be ready if they watch you eating, reach for your food, and seem eager to be fed.
- **Can he move food from a spoon into his throat?** If you offer a spoon of rice cereal and he pushes it out of his mouth and it dribbles onto his chin, he may not have the ability to move it to the back of his mouth to swallow it. It's normal. Remember, he's never had anything thicker than breast milk or formula before, and this may take some getting used to. Try diluting it the first few times, then gradually thicken the texture. You may also want to wait a week or two and try again.
- **Is he big enough?** Generally, when infants double their birth weight (typically at about 4 months) and weigh about 13 pounds or more, they may be ready for solid foods.

NOTE: The AAP recommends breastfeeding as the sole source of nutrition for your baby for about 6 months. When you add solid foods to your baby's diet, continue breastfeeding until at least 12 months. You can continue to breastfeed after 12 months if you and your baby desire.

Check with your child's doctor about vitamin D and iron supplements during the first year.

How do I feed my baby?

Start with half a spoonful or less and talk to your baby through the process ("Mmm, see how good this is?"). Your baby may not know what to do at first. She may look confused, wrinkle her nose, roll the food around her mouth, or reject it altogether.

One way to make eating solids for the first time easier is to give your baby a little breast milk and/or formula first, then switch to very small half-spoonfuls of food, and finish with more breast milk and/or formula. This will prevent your baby from getting frustrated when she is very hungry.

Do not be surprised if most of the first few solid-food feedings wind up on your baby's face, hands, and bib. Increase the amount of food gradually, with just a teaspoonful or two to start. This allows your baby time to learn how to swallow solids.

Do not make your baby eat if she cries or turns away when you feed her. Go back to nursing or bottle-feeding exclusively for a time before trying again. Remember that starting solid foods is a gradual process and at first your baby will still be getting most of her nutrition from breast milk and/or formula.

NOTE: Do not put baby cereal in a bottle because your baby could choke. It also may increase the amount of food your baby eats and can cause your baby to gain too much weight. However, cereal in a bottle may be recommended if your baby has reflux. Check with your child's doctor.

Which food should I give my baby first?

For most babies it does not matter what the first solid foods are. By tradition, single-grain cereals are usually introduced first. However, there is no medical evidence that introducing solid foods in any particular order has an advantage for your baby. Though many pediatricians will recommend starting vegetables before fruits, there is no evidence that your baby will develop a dislike for vegetables if fruit is given first. Babies are born with a preference for sweets, and the order of introducing foods does not change this. If your baby has been mostly breastfeeding, he may benefit from baby food made with meat, which contains more easily absorbed sources of iron and zinc that are needed by 4 to 6 months of age. Check with your child's doctor.

Baby cereals are available premixed in individual containers or dry, to which you can add breast milk, formula, or water. Whichever type of cereal you use, make sure that it is made for babies and iron-fortified.

When can my baby try other food?

Once your baby learns to eat one food, gradually give him other foods. Give your baby one new food at a time, and wait at least 2 to 3 days before starting another. After each new food, watch for any allergic reactions such as diarrhea, rash, or vomiting. If any of these occur, stop using the new food and consult with your child's doctor.

Generally, meats and vegetables contain more nutrients per serving than fruits or cereals. Many pediatricians recommend against giving eggs and fish in the first year of life because of allergic reactions, but there is no evidence that introducing these nutrient-dense foods after 4 to 6 months of age determines whether your baby will be allergic to them.

Within a few months of starting solid foods, your baby's daily diet should include a variety of foods each day that may include the following: breast milk and/or formula, meats, cereal, vegetables, fruits, eggs, and fish.

NOTE: If you make your own baby food, be aware that home-prepared spinach, beets, green beans, squash, and carrots are not good choices during early infancy. They may contain large amounts of nitrates. Nitrates are chemicals that can cause an unusual type of anemia (low blood count) in young babies. Commercially prepared vegetables are safer because the manufacturers test for nitrates. Peas, corn, and sweet potatoes are better choices for home-prepared baby foods.

When can I give my baby finger foods?

Once your baby can sit up and bring her hands or other objects to her mouth, you can give her finger foods to help her learn to feed herself. To avoid choking, make sure anything you give your baby is soft, easy to swallow, and cut into small pieces. Some examples include small pieces of banana, wafer-type cookies, or crackers; scrambled eggs; well-cooked pasta; well-cooked chicken finely chopped; and well-cooked and cut up yellow squash, peas, and potatoes.

At each of your baby's daily meals, she should be eating about 4 ounces, or the amount in one small jar of strained baby food. Limit giving your baby foods that are made for adults. These foods often contain more salt and other preservatives.

If you want to give your baby fresh food, use a blender or food processor, or just mash softer foods with a fork. All fresh foods should be cooked with no added salt or seasoning. Though you can feed your baby raw bananas (mashed), most other fruits and vegetables should be cooked until they are soft. Refrigerate any food you do not use, and look for any signs of spoilage before giving it to your baby. Fresh foods are not bacteria-free, so they will spoil more quickly than food from a can or jar.

NOTE: Do not give your baby any food that requires chewing at this age. Do not give your baby any food that can be choking hazards, including hot dogs (including meat sticks [baby food "hot dogs"]); nuts and seeds; chunks of meat or cheese; whole grapes; popcorn; chunks of peanut butter; raw vegetables; fruit chunks, such as apple chunks; and hard, gooey, or sticky candy.

What changes can I expect after my baby starts solids?

When your baby starts eating solid foods, his stools will become more solid and variable in color. Because of the added sugars and fats, they will have a much stronger odor too. Peas and other green vegetables may turn the stool a deep-green color; beets may make it red. (Beets sometimes make urine red as well.) If your baby's meals are not strained, his stools may contain undigested pieces of food, especially hulls of peas or corn, and the skin of tomatoes or other vegetables. All of this is normal. Your baby's digestive system is still immature and needs time before it can fully process these new foods. If the stools are extremely loose, watery, or full of mucus, however, it may mean the digestive tract is irritated. In this case, reduce the amount of solids and introduce them more slowly. If the stools continue to be loose, watery, or full of mucus, consult your child's doctor to find the reason.

Should I give my baby juice?

Babies do not need juice. Babies younger than 6 months should not be given juice. However, if you choose to give your baby juice, do so only after 6 months of age, give only 100% fruit juice, and offer it only in a cup, not in a bottle. To help prevent tooth decay, do not put your child to bed with a bottle. If you do, make sure it contains only water.

Limit juice to no more than 4 ounces a day and offer it only with a meal or snack. Any more than this will reduce her appetite for other, more nutritious foods, including breast milk and/or formula. Too much juice also can cause diaper rash, diarrhea, or excessive weight gain.

Using a high chair

The following are safety tips when using a high chair:

- Make sure the high chair you use cannot be tipped over easily.
- If the chair folds, be sure it is locked each time you set it up.
- Whenever your child sits in the chair, use the safety straps, including the crotch strap. This will prevent your child from slipping down, which could cause serious injury or even death. Never allow your child to stand in the high chair.
- Do not place the high chair near a counter or table. Your child may be able to push hard enough against these surfaces to tip the chair over.
- Never leave a young child alone in a high chair, and do not allow older children to climb or play on it because this could also tip it over.
- A high chair that hooks on to a table is not a good substitute for a freestanding one. If you plan to use this type of chair when you eat out or travel, look for one that locks on to the table. Be sure the table is heavy enough to support your child's weight without tipping. Also, check to see whether your child's feet can touch a table support. If your child pushes against the table, it may dislodge the seat.

Does my baby need water?

Healthy babies do not need extra water. Breast milk and/or formula provides all the fluids they need. However, with the introduction of solid foods, water can be added to your baby's diet. Also, a small amount of water may be needed in very hot weather, but check with your child's doctor about how much is safe. And if you live in an area where the water is fluoridated, drinking water also will help prevent future tooth decay.

Good eating habits start early

It is important for your baby to get used to the process of eating—sitting up, taking food from a spoon, resting between bites, and stopping when full. These early experiences will help your child learn good eating habits throughout life.

Encourage family meals from the first feeding. When you can, the whole family should eat together. Research suggests that having dinner together as a family on a regular basis has positive effects on the development of children.

Remember to offer a good variety of healthy foods that are rich in the nutrients your child needs. Watch your child for cues that he has had enough to eat. Do not overfeed!

If you have any questions about your child's nutrition, including concerns about your child eating too much or too little, talk with your child's doctor.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

American Academy
of Pediatrics




DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
Web site—www.healthychildren.org

Copyright © 2008
American Academy of Pediatrics, Updated 2/12
All rights reserved.



Baby-sitting Reminders

IMPORTANT PHONE NUMBERS

PARENTS _____ POLICE _____
NEIGHBOR(S) _____ POISON CENTER _____
DOCTOR _____ HOME PHONE _____
FIRE/RESCUE _____ HOME ADDRESS _____

PARENTS SHOULD

- Meet the sitter and check references and training in advance.
- Be certain the sitter has had first aid training and knows CPR.
- Be sure the sitter is at least 13 years old and mature enough to handle common emergencies.
- Have the sitter spend time with you before baby-sitting to meet the children and learn their routines.
- Show the sitter around the house. Point out fire escape routes and potential problem areas. Instruct the sitter to leave the house right away in case of fire and to call the fire department from a neighbor's house.
- Discuss feeding, bathing, and sleeping arrangements for your children.
- Tell your sitter of any allergies or specific needs your children have.
- Have emergency supplies available including a flashlight, first aid chart, and first aid supplies.
- Tell the sitter where you will be and when you will return.
- Be sure any guns are stored unloaded in a locked cabinet, and lock and store the ammunition in a separate place.

SITTERS SHOULD

- Be prepared for an emergency.
- Always phone for help if there are any problems or questions.
- Never open the door to strangers.
- Never leave the children alone in the house — even for a minute.
- Never give the children any medicine or food unless instructed to do so by the parents.
- Remember that your job is to care for the children. Tender loving care usually quiets an unhappy child.

From Your Doctor
The Perkins Pediatric Clinic, LLC
Michael W. Perkins, MD
614 North Pine Street
DeRidder, LA 70634
(337) 462-6000
www.perkinspediatrics.com


The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on the individual facts and circumstances.

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Teething in Children: Care Instructions



		Age tooth comes in	Age tooth falls out
Upper teeth	Central incisor	8-12 mos.	6-7 yrs.
	Lateral incisor	9-13 mos.	7-8 yrs.
	Canine (cuspid)	16-22 mos.	10-12 yrs.
	First molar	13-19 mos.	9-11 yrs.
	Second molar	25-33 mos.	10-12 yrs.
Lower teeth	Second molar	23-31 mos.	10-12 yrs.
	First molar	14-18 mos.	9-11 yrs.
	Canine (cuspid)	17-23 mos.	9-12 yrs.
	Lateral incisor	10-16 mos.	7-8 yrs.
	Central incisor	6-10 mos.	6-7 yrs.

© Healthwise, Incorporated

Your Care Instructions

Teething is the normal process in which your baby's first set of teeth (primary teeth) break through the gums (erupt). Teething usually begins at around 6 months of age, but it is different for each child. Some children begin teething at 3 to 4 months, while others do not start until age 12 months or later. A total of 20 teeth erupt by the time a child is about 3 years old. Usually teeth appear first in the front of the mouth. Lower teeth usually erupt 1 to 2 months earlier than their matching upper teeth. Girls' teeth often erupt sooner than boys' teeth.

Your child may be irritable and uncomfortable from the swelling and tenderness at the site of the erupting tooth. These symptoms usually begin about 3 to 5 days before a tooth erupts and then go away as soon as it breaks the skin. Your child may bite on fingers or toys to help relieve the pressure in the gums. He or she may refuse to eat and drink because of mouth soreness. Children sometimes drool more during this time. The drool may cause a rash on the chin, face, or chest.

Teething may cause a mild increase in your child's temperature. But if the temperature is higher than 100.4 F (38 C), look for symptoms that may be related to an infection or illness.

You might be able to ease your child's pain by rubbing the gums and giving your child safe objects to chew on.

Follow-up care is a key part of your child's treatment and safety. Be sure to make and go to all appointments, and call your doctor if your child is having problems. It's also a good idea to know your child's test results and keep a list of the medicines your child takes.

How can you care for your child at home?

- Give acetaminophen (Tylenol) or ibuprofen (Advil, Motrin) for pain or fussiness. Read and follow all instructions on the label.

- Gently rub your child's gum where the tooth is erupting for about 2 minutes at a time. Make sure your finger is clean, or use a clean teething ring.
- Do not use teething gels for children younger than age 2. Ask your doctor before using mouth-numbing medicine for children older than age 2. The U.S. Food and Drug Administration (FDA) warns that some of these can be dangerous. Talk to your child's doctor about other teething remedies.
- Give your child safe objects to chew on, such as teething rings. Do not use fluid-filled teethingers.
- If your child is eating solids, try offering cold foods and fluids, which help to ease gum pain. You can also dip a clean washcloth in water, freeze it, and let your child chew on it.

When should you call for help?



Call your doctor now or seek immediate medical care if:

- Your child has a fever.
- Your child keeps pulling on his or her ears.
- Your child has diarrhea or a severe diaper rash.

Watch closely for changes in your child's health, and be sure to contact your doctor if:

- You think your child has tooth decay.
- Your child is 18 months old and has not had an erupting tooth yet.

Where can you learn more?

Go to <https://www.healthwise.net/patiented>

Enter **C015** in the search box to learn more about "**Teething in Children: Care Instructions**".

©2006-2019 Healthwise, Incorporated.

Care instructions adapted under license by The Perkins Pediatric Clinic, LLC. This care instruction is for use with your licensed healthcare professional. If you have questions about a medical condition or this instruction, always ask your healthcare professional. Healthwise, Incorporated disclaims any warranty or liability for your use of this information.