



# Bright Futures Parent Handout

## 15 to 17 Year Visits

Here are some suggestions from Bright Futures experts that may be of value to your family.

PHYSICAL GROWTH AND DEVELOPMENT

### Your Growing and Changing Teen

- Help your teen visit the dentist at least twice a year.
- Encourage your teen to protect her hearing at work, home, and concerts.
- Keep a variety of healthy foods at home.
- Help your teen get enough calcium.
- Encourage 1 hour of vigorous physical activity a day.
- Praise your teen when he does something well, not just when he looks good.

RISK REDUCTION

### Healthy Behavior Choices

- Talk with your teen about your values and your expectations on drinking, drug use, tobacco use, driving, and sex.
- Be there for your teen when she needs support or help in making healthy decision about her sexual behavior.
- Support safe activities at school and in the community.
- Praise your teen for healthy decisions about sex, tobacco, alcohol, and other drugs.

VIOLENCE AND INJURY PREVENTION

### Violence and Injuries

- Do not tolerate drinking and driving.
- Insist that seat belts be used by everyone.
- Set expectations for safe driving.
  - Limit the number of friends in the car, nighttime driving, and distractions.
- Never allow physical harm of yourself, your teen, or others at home or school.
- Remove guns from your home. If you must keep a gun in your home, make sure it is unloaded and locked with ammunition locked in a separate place.
- Teach your teen how to deal with conflict without using violence.
- Make sure your teen understands that healthy dating relationships are built on respect and that saying "no" is OK.

EMOTIONAL WELL-BEING

### Feelings and Family

- Set aside time to be with your teen and really listen to his hopes and concerns.
- Support your teen as he figures out ways to deal with stress.
- Support your teen in solving problems and making decisions.
- If you are concerned that your teen is sad, depressed, nervous, irritable, hopeless, or angry, talk with me.

SOCIAL AND ACADEMIC COMPETENCE

### School and Friends

- Praise positive efforts and success in school and other activities.
- Encourage reading.
- Help your teen find new activities she enjoys.
- Encourage your teen to help others in the community.
- Help your teen find and be a part of positive after-school activities and sports.
- Encourage healthy friendships and fun, safe things to do with friends.
- Know your teen's friends and their parents, where your teen is, and what he is doing at all times.
- Check in with your teen's teacher about her grades on tests.
  - Attend back-to-school events if possible.
  - Attend parent-teacher conferences if possible.



## American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of *Bright Futures Tool and Resource Kit*. Copyright © 2010 American Academy of Pediatrics. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.



Illustration by Billy Nuñez, age 16

# TEEN DATING VIOLENCE: TIPS FOR PARENTS

## PARENTS NEED TO KNOW WHAT'S GOING ON

- Learn the myths and facts about dating violence.
- Some teens who are abused don't talk about it.
- There are warning signs of abuse.
- Teach your teenager that respect is the most important thing in a dating relationship.
- Learn how to talk with your child about dating violence.

**R**omantic and sexual feelings develop during the teen years. Teenagers are starting to date and experiment with different types of relationships. It is fun and exciting to meet someone new, and sad and difficult to break up.

As a parent, you can help your teenager make good decisions about dating. With guidance and support, teens can learn about healthy relationships and get the strength and courage needed to leave those that are not.

**There is no place for verbal or physical abuse in a healthy and respectful relationship.**

## MYTHS AND FACTS

MYTH: "Teen dating violence is just another way of saying rape."

FACT: Abuse comes in many forms. Besides sexual violence, it also includes:

- Yelling, swearing, put-downs, and threats
- Being pushed around or hit
- Controlling, bossy, and bullying behavior

MYTH: "Oh, it's not that common."

FACTS:

- More than 1 in 10 teenagers experience physical violence in a dating relationship.
- When threats and emotional abuse are included, it's even higher.

MYTH: "It only happens to kids from bad homes."

FACT: Dating violence is not limited to families with a history of violence. It happens to teens from families of all cultures, income levels, and educational backgrounds.

MYTH: "It can't happen to my child."

FACTS:

- Boys, as well as girls, can be victims of dating violence.
- It can happen in any type of relationship—straight, gay, or lesbian.
- It can occur at any time in a relationship—those just starting or ones that have been going on for a while.

## WHY TEENS ARE SILENT

There are many reasons why teens don't tell their parents or friends about the violence they are experiencing. They may:

- Feel embarrassed, ashamed, or afraid of getting hurt.
- Be convinced it is their fault and do not know that it is abuse.
- Feel afraid they will be forced to break up or of losing privileges.
- Feel certain that being in a relationship is the most important thing in their life.
- Confuse jealousy with love, and think this is how it should be.



- Be afraid of losing friends. They may think friends would not believe this is happening, and they may feel alone after having lost touch with friends.
- Know the abuser will act nice some of the time, and they are happy when this happens.
- Feel hopeful that things will get better and convinced that they can help or change their partner.

## **WARNING SIGNS OF ABUSE**

Some of the following may be just part of being a teenager. But, when these changes happen suddenly or without explanation, there is cause for concern.

### **School**

- Failing grades
- Dropping out of school or school activities

### **Attitude/mood**

- Difficulty making decisions
- Changes in personality, becoming anxious or depressed
- Acting out or being secretive
- Avoiding eye contact
- Having “crying jags” or getting “hysterical”
- Constantly thinking about dating partner

### **Physical appearance**

- Bruises, scratches, or other injuries
- Sudden changes in clothes or make-up

### **Activities**

- Avoiding friends or changing peer groups
- Giving up activities, interests, or family time that previously had been important
- Changes in eating or sleeping habits
- Using alcohol, tobacco, or other drugs

### **Pregnancy**

- Some teenagers believe that having a baby will help make things better.
- Some girls are forced to have sex.

## **ABUSIVE BEHAVIOR**

It is important for you to recognize signs of an abusive relationship.

- Wanting to get serious quickly and refusing to take NO for an answer
- Acting jealous and possessive and wanting to pick partner’s friends and activities
- Being controlling and bossy and making all the decisions, not taking opinions of others seriously, and always checking up on partner
- Using threats and “put-downs” when alone or with friends
- Using guilt trips like, “If you really loved me, you would\_\_\_\_\_.”
- Blaming the victim for what is wrong, like saying, “It’s because of you that I get so mad.”
- Apologizing or giving excuses for violent behavior like, “I promise I’ll never do it again,” or “I was drinking and just didn’t know that I \_\_\_\_\_.”

**Your child needs help in either case—  
victim or abuser**

## **TIPS FOR PARENTS**

### **Talk about healthy dating relationships and possible problems.**

It is never too early or too late to teach about respect. Respect for self and others is important in any relationship.

### **If you think your teenager already may be involved with an abusive partner:**

- Give your teen a chance to talk. Listen quietly to the whole story.
- Tell your child that you are there to help, not to judge.
- If your teen does not want to talk with you, help find another trusted person for your child to talk with.
- Focus on your child’s safety and self-esteem. Point out how unhappy your teenager seems to be while with this person and the possibility of danger or harm. Do not “put down” the abusive partner.
- Let your child know that abuse always gets worse. What may start as minor verbal or physical abuse is very likely to get worse if not stopped immediately.

## If your teenager tries to break up with an abusive partner:

- Advise that the breakup be definite and final.
- Develop a safety plan with your teen ahead of time.
- Support your teenager's decision and be ready to help.
- Make sure your teen takes all necessary safety measures.
  - Avoid walking or riding alone, and always travel with another friend.
  - Consider changing class schedules.
  - Get help from the guidance counselor, school principal, or if necessary, the police.

## WHAT YOU CAN SAY

It is important for your teen to talk about dating violence, either with you or with another trusted adult. Many people who have been victims of dating violence have been able to change their lives after they began talking with others.

- "I care about what happens to you. I love you and I want to help."
- "If you feel afraid, it may be abuse. Sometimes people act in ways that are scary and make you feel threatened, even without using physical violence. Pay attention to what you are feeling."
- "No one has the right to hit you. No one has the right to control you or what you do."
- "The abuse is not your fault. You are not to blame, no matter how guilty \_\_\_\_\_ is trying to make you feel. This should not be happening to you."
- "It is \_\_\_\_\_ who has a problem, not you. It is not your responsibility to help \_\_\_\_\_ change."

**Ask for teen dating violence prevention and intervention programs at your teenager's school or through religious or other community groups.**

The Perkins Pediatric Clinic, LLC  
Michael W. Perkins, MD  
614 North Pine Street  
DeRidder, LA 70634

phone: (337) 462-6000  
[www.perkinspediatrics.com](http://www.perkinspediatrics.com)

Connected Kids are Safe, Strong, and Secure

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

The American Academy of Pediatrics is an organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

This project was supported by Grant No. 2001-JN-FX-0011 awarded by the Office of Juvenile Justice and Delinquency Prevention, Office of Justice Programs, U.S. Department of Justice. Points of view or opinions in this document are those of the author and do not necessarily represent the official position or policies of the U.S. Department of Justice. Adapted with permission from the Massachusetts Medical Society's "Teen Dating Violence" Parent Education Card, Copyright 2000.

All rights reserved.  
HE50400

Graphic design and illustrations by Artists For Humanity, a non profit arts and entrepreneurship program for Boston teens.

American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

# STDs/STIs: Protecting Yourself

What is an STD?

An STD -- short for sexually transmitted disease -- is an infection you can get by having sex. Sex in this case includes intercourse, anal sex, oral sex or skin-to-skin contact. Some of the most common STDs are listed below.

How can I PROTECT myself against an STD?

The only 100% sure way to protect yourself is NOT TO HAVE SEX. It only takes one sexual contact with someone who has an STD to catch it. If you are sexually active, protect yourself by using a latex condom and a spermicide foam, cream or jelly with nonoxynol-9 (check the label on the box) every time you have sex.

Did you know that... **ONE OUT OF EVERY 8 TEENS GET AN STD.**

**Pubic lice** (crabs): Girls and guys may have redness and itching around the genitals.

**Trichomoniasis:** Girls can have a heavy, greenish-yellow frothy discharge and pain when urinating. Guys don't usually have symptoms.

**Chlamydia:** Girls may have no symptoms or may have pain when urinating, itching around the vagina, yellow fluid (discharge) from the vagina, bleeding between periods or pain in the lower abdomen. Guys may feel a burning when urinating and have milky colored discharge from the penis. If not treated, chlamydia can cause infertility and other problems in girls and painful swelling of the scrotum in guys.

**Syphilis:** Symptoms: An early symptom is a red PAINLESS sore, called a chancre. The sore can be on the penis, vagina, tongue or throat. The glands near the sore may be swollen. After a few months, both guys and girls can get a fever, sore throat, headache, or pain in their joints. Another symptom is a scaly rash on the palms of the hands or the bottom of the feet. The sores and other symptoms go away, but this does not mean that the infection is gone. Syphilis can cause serious health problems if it's not treated.

**HIV/AIDS:** HIV (human immunodeficiency virus) causes AIDS. HIV makes the body's immune system weak so it can't fight disease. Symptoms may take years to develop, and can include infections, feeling tired for no reason, and night sweats. HIV CANNOT BE CURED.

**Herpes:** Girls and guys may have tingling, PAIN or itching around the vagina or penis. Small blisters can form in these areas and then break open. When they break open, the sores can cause a burning feeling. It may hurt to urinate. Some people have swollen glands, fever and body aches. The sores and other symptoms go away, but this does not mean that the infection is gone. The sores and blisters can come back (called an "outbreak").  
HERPES CANNOT BE CURED.

**Gonorrhea:** Girls may have no symptoms or may have white, green or yellow discharge from the vagina, pain when urinating, bleeding between periods, heavy bleeding during a period or a fever. Both girls and guys can get sore throats if they've had mouth to penis or vagina contact (oral sex). Guys may have thick, yellow discharge from the penis and pain when urinating. The opening of the penis may be sore.

**HPV/Genital Warts:** HPV (human papillomavirus) can cause warts in or around the vagina, penis or rectum. In girls, the warts can be inside the body on the cervix or vagina so you can't see them. Or they can be on the outside of the body but be too small to see. The warts usually don't hurt. HPV CANNOT BE CURED.

**How do I know if my PARTNER has an STD?**

Ask. Although it may be uncomfortable, talk to your partner before having any sexual contact. Ask if he or she is at risk for having an STD. Some of the risk factors are having sex with several partners, using injected drugs, and having had an STD in the past. To be safe, protect yourself no matter what the person says. You must also tell your partner if you have an STD. You aren't doing yourself or your partner any favors by trying to hide it.

**How do I know if I have an STD?**

Watch for the symptoms listed above, but remember that most STDs don't cause any symptoms.

If you are sexually active, you should get regular check-ups and tell your doctor that you are sexually active. If you're worried that your parents will find out, you can ask your doctor not to discuss it with your parents. You can also go to a free health clinic. Don't let fear keep you from getting checked out. Imagine how you'd feel -- and how your parents would feel -- if you got really sick because you didn't get help. If you find out you have an STD, both you and your partner should get treated right away.

**Can STDs be CURED?**

Some can. STDs like chlamydia that are caused by bacteria can be cured with antibiotics. But STDs caused by a virus (like HIV or herpes) can't be cured. Your doctor can only treat the symptoms that the virus causes.

Don't wait to be treated! Early treatment helps prevent serious health problems. Even if medicine can't completely cure the STD, it can help keep you from getting sick. If you are given medicine for an STD, take it exactly as the doctor says.

**For more information:**

**National STD Hotline:**

800-227-8922

**National AIDS Hotline:**

800-342-2437 (English)

800-344-7432 (Spanish)

800-243-7889 (Hearing Impaired)

# Effects of Premarital Sexual

## Relations

### WHAT'S THE BIG DEAL:

Lucy and Ricky Ricardo sleep in separate twin beds in the 1950s comedy "I Love Lucy" to avoid any innuendoes; now media rages with suggestive material. Nick Lebel of Focus Adolescent Services reports:

- In 2003, 83% of the episodes of the top 20 shows among teen viewers contained some sexual content, including 20% with sexual intercourse.
- 42% of the songs on the top CDs in 1999 contained sexual content. 19% included direct descriptions of sexual intercourse.
- On average, music videos contain 93 sexual situations per hour, including eleven "hard core" scenes depicting behaviors such as intercourse and oral sex.
- Before parents raised an outcry, Abercrombie and Fitch marketed a line of thong underpants decorated with sexually provocative phrases such as "Wink Wink" and "Eye Candy" to 10-year-olds.
- Neilson estimates that 6.6 million children ages 2-11 and 7.3 million teens ages 12-17 watched Justin Timberlake rip open Janet Jackson's bodice during the 2004 Super Bowl halftime show.

The internet is also a source of the perversion of sexuality; there are more than one million porn sites, and most of them are easy to find, even by accident. Porn, particularly, teaches teens to view women as mere sex objects, and later, porn becomes part of their sexuality. Paul Schenk, Psy.D. says, "It has the potential for teenagers to really mess up what their attitudes are about sexuality and really mess up relationships." "Although most teenage girls believe that sex equals love, other teens-especially boys-believe that sex is not the ultimate expression of the ultimate commitment, but a casual activity". Why shouldn't they? Is this not the message television is sending? Very seldom, if ever, does the entertainment business depict the sexual risks such as incurable diseases and unwanted pregnancies.

In the controversy of premarital sex, one side may argue it produces positive outcomes. The youth are forced to accept responsibilities such as using contraceptives, caring for another life, or dealing with the risks and consequences. In various cases, the teens use the situation as a turning point in life and choose to settle down and act more maturely. To ease the uneasiness adolescents sometimes feel when dealing with the opposite sex, some advocates proclaim it is beneficial to "test the waters" and to acquire experience. In this growing epidemic of sexual relations among teens, the bad results by far outweigh the good.

In his thesis on "The Phenomenon of Premarital Sex," Vic Fabe relates the information he discovered, "In the 1960s 25 percent of young men and 45 percent of young women were virgins at the age of nineteen; by the 1980s, fewer than 20 percent of males and females were." A survey performed in 2003 by a psychology unit found about one in five teens report having sex before age fifteen. More than half of fifteen to seventeen-year-olds have been with someone in a

sexual way. Other surveys have found that nearly two in three teens will have had sex by the time they graduate from high school.

The words most commonly uttered by teens choosing to become sexually promiscuous tend to relate to pregnancy: "As long as she doesn't get pregnant" or "If I get pregnant my parents will shoot me." Out-of-wedlock children constitute one in three of all births. In the United States in 1994, twenty-two percent of the births were to single girls eighteen years old or younger. Statistics show the teen pregnancy rate has fallen steadily between 1990 and 2000; however, the numbers are still quite high. In 2004 more than 415,000 babies were born to teens, and more than eighty percent of these births were to unmarried teens. These numbers do not include the babies conceived and miscarried or aborted.

Pregnancy presents many difficult decisions for the teen mother and father: Abortion? Adoption? Keep? "When teenagers become pregnant, abortion is a likely result. For girls under 15, there are 8 abortions to every 10 live births. For girls 15-19, there are 4 abortions to every 10 live births." If against terminating the pregnancy, adoption is another option, but for most females it is in their intricate nature to be unable to part with the infant. In deciding to keep the child, the mother will face many challenges ahead. More than three-fifths of teen mothers are poverty-stricken at the time of their child's birth and over four-fifths eventually live below the poverty level. Teen childbearing costs U.S. taxpayers an estimated \$7 billion per year. Low education levels account for a major portion of government dependency of teenagers; only about fifty percent of teen mothers are likely to finish high school. Teen mothers have a higher percentage likelihood of experiencing stressful major life events such as having a small child die or taken away, a disabling or life-threatening accident or injury, sexual assault or abuse, or an immediate family member addicted to drugs and alcohol. Despite these details, pregnancy should be the least of teenagers' worries.

For every unwed teenager who gets pregnant this year, 10 teenagers will get an STD. Approximately 333 million new cases of Sexually Transmitted Diseases are reported in the world each year. Teens are only ten percent of the population, but they account for twenty-five percent of all STDs. One out of four Americans has an incurable, viral STD; fifty percent of Americans age fourteen to twenty-five have an incurable, viral STD.

The World is plagued with over thirty-nine Sexually Transmitted Diseases. Hepatitis B is an extremely prevalent STD, and it is the biggest battle for pediatricians; however, the STD that arouses the biggest fear in society is HIV which eventually progresses into AIDS. Approximately twenty-five percent of HIV transmission in the United States is estimated to occur among people younger than twenty-one years of age. The human immunodeficiency virus destroys a certain kind of blood cell which is crucial to the normal function of the immune system. The symptoms of HIV are similar to those of many other diseases, so they cannot be relied on to determine HIV infection. Warning signs of the human immunodeficiency virus are as follows:

Rapid weight loss, dry cough, recurring fever or profuse night sweats, profound and unexplained fatigue, swollen lymph glands in the armpits, groin or neck, diarrhea that lasts for more than a week, white spots or unusual blemishes on the tongue, in the mouth or in the throat, pneumonia, red, brown, pink or purplish blotches on or under the skin or inside the mouth, nose or eyelids, memory loss, depression and other neurological disorders.

Many carriers of HIV do not have any symptoms at all for years. Many people contract HIV as teens but do not develop AIDS until later on in life. This fact is why comparatively few teenagers die of Acquired Immunodeficiency Syndrome. An estimated one-half million people in the United States died with AIDS in 2002, while only about two thousand of these were adolescents.

On average, the first stage of syphilis occurs twenty-one days after onset of infection. Anytime between ten to ninety days, the person may experience a firm, round, small, and painless sore where the syphilis entered the body. It lasts from one to six weeks, and it heals without treatment. During the second stage, the person may have a red or brownish head-to-toe rash, fever, swollen lymph glands, sore throat, patchy hair loss, headaches, weight loss, muscle aches, and fatigue. Once again, the symptoms of secondary syphilis will go away without treatment; however, without treatment the infection will progress to the late stages of disease. The third stage, also referred to as the hidden stage,



begins when the second disappears. The infection remains in the body even though there are no symptoms. At this stage there is no treatment. It may damage internal organs including: the brain, nerves, eyes, heart, blood vessels, liver, bones, and joints. Signs consist of difficulty with muscle movements, paralysis, numbness, gradual blindness, and dementia.

Between twenty-eight and forty-six percent of men and women under the age of twenty-five are infected with human papillomavirus. The virus lives in the skin or mucus membranes and usually causes no symptoms. It infects the genitalia, causes warts, and, in some instances, leads to anal or genital cancers.

Among the other common STDs are Chlamydia, more so among teenagers than older men and women, and Gonorrhea, which cause painful discharges and may lead to sterility and to pelvic inflammatory disease in females; herpes, which causes genital ulcers; and trichomoniasis, which causes a discharge and itching in females and advances to urethritis in males.

Among teens who have not yet had sex, nearly a third say they have been "intimate" with a partner. So if teenagers decide not to have intercourse, then how far is too far? Is oral sex really sex? As one eighth-grader put it, "What's the big deal? President Clinton did it." The Missouri Division of Family Services of Focus Adolescent Services reports their categorization of the "normal range" of sexual behavior for teens:

- Sexually explicit conversations with peers
- Obscenities and jokes within cultural norm
- Sexual innuendo, flirting and courtship
- Interest in erotica
- Solitary masturbation
- Hugging, kissing, holding hands
- Foreplay, (petting, making out, fondling) and mutual masturbation: Moral, social or familial rules may restrict, but these behaviors are not abnormal, developmentally harmful, or illegal when private, consensual, equal and non-coercive.
- Monogamist intercourse: Stable monogamy is defined as a single sexual partner throughout adolescence. Serial monogamy indicates long-term (several months or years) involvement with a single partner which ends and is then followed by another.

Alicia in Pure Excitement offers her view on the topic, "Now I know that 'too far' doesn't mean only intercourse, but also the stages leading up to it. Too far is when sexual thoughts take over your relationship. Too far is when you don't want to stop." Oral sex, like other methods of sex, carries with it the risk of serious, untreatable and even life-threatening diseases in both young men and women. Oral sex has been found to spread HIV, HPV, syphilis, gonorrhea, chlamydia, genital herpes, and possibly hepatitis C. Studies have shown that petting also leads to the transfer of some STDs. Anywhere genital contact is involved, a person put himself or herself at risk.

Society and even the teenagers themselves focus on the physical aspects of the issue, and they neglect to address the emotional tolls premarital sex has on the younger generation. The sexual relationship is predominantly emotional and moral rather than physical in character. All actions, decisions, and attitudes begin in the mind. The mind is susceptible to manipulation by almost any attractive outside source. The brain has ten billion cells that act like tiny storage building that capture every voluntary and involuntary sight and sound a person ever sees or hears, especially when connected with music. The subconscious mind absorbs thoughts even when the conscious mind does not command it to do so. The mind is the foundation of human emotion, and it plays a large role in the aftermath of a premarital sexual experience. Premarital sex often leads to feelings of fear, guilt, regret, embarrassment, tension, distrust, lack of respect, resentment, confusion, and so much more.

Amidst all the others, depression is a prime mentality scientist have linked with premarital sex among teenagers. The Heritage Foundation found when compared to teens who are not sexually active, teenage boys and girls who are sexually active are significantly:

**Less likely to be happy and more likely to feel DEPRESSED.**

**More likely to attempt SUICIDE.**

In an evaluation 25.3 percent of sexually active teenage girls rate themselves as being depressed all, most, or a lot of the time. Only 7.7 percent of non-sexually active teenage girls report feeling depressed at the same level. While the numbers for teenage boys are much lower, the distinction between sexually active and not sexually active is apparent. 8.3 percent of sexually active teenage boys report being depressed all, most, or a lot of the time. Only 3.4 percent of non-sexually active teenage boys describe themselves in the same way. 14.3 percent of girls who are sexually active report having attempted suicide. Only 5.1 percent of sexually abstinent girls have attempted suicide. Six percent of boys who are sexually active have attempted suicide. Only 0.7 percent of sexually abstinent boys have attempted suicide.

**Sexually active GIRLS are more than three times more likely to be depressed.**

**Sexually active GIRLS are nearly three times more likely to attempt suicide.**

**Sexually active BOYS are more than twice as likely to be depressed.**

**Sexually active BOYS are eight times more likely to attempt suicide.**

Most sexually active teens express reservations and concerns about their personal sexual experiences. Most boys and virtually three-fourths of girls regard their own initial sexual experience adversely—"as an event they wish they had avoided." Nearly two out of three of all teens who have already had sex wish they had waited longer before engaging in sex. Sixty-two percent of teens regret ever getting started. Janet shares her remorse, "The thing I regret most in my life would have to be losing my virginity. I was so young, and most people don't think 12-year-olds (7th grade) even know about sex." Jason gives his testimony in Pure Excitement, "I have made many mistakes in my life, including having had sex once. Afterward, I felt very bad and empty inside."

A shocking percentage of Americans condone premarital sexual relations; forty-one percent of the population thinks it is "not wrong at all." Merely twenty-nine percent think it is "always wrong." Twelve percent think that it is wrong with a few exceptions, and eighteen percent think that it is right with a few exceptions. "The average person 'falls in love' about seven times before he or she gets married. And there are those who say sex is okay if you're 'in love.' The problem with that is that your bride or bridegroom isn't supposed to be your eighth honeymoon."

Sex outside of marriage also has far-reaching psychological effects. Partakers usually obtain a fear of commitment which follows them later in life. They are less likely to have a satisfying marriage relationship, and non-virgins increase their odds of divorce by about sixty percent. "The Seven Effects of Premarital Sex" summarizes the extensive effects of premarital relations on a future marriage:


1. Premarital sex tends to break up couples before marriage takes place.
2. Many men do not want to marry a woman who has had intercourse with someone else. The strange logic seems to be, "Its (sic) okay for me to have sex with the girl you marry, but it's not okay for you to have sex with the girl I marry."

3. Those who have premarital sex tend to have less happy marriages. The physical relationship is an inadequate foundation upon which to build a lasting relationship.
4. Those who have premarital sex are more likely to have their marriages end in divorce.
5. Person and couples who have had premarital sex are more likely to have extramarital affairs as well. This is especially true for women; those who engaged in sex before marriage are more than twice as likely to have extramarital affairs as those who did not have premarital sex.
6. Having premarital sex may fool you into marrying a person who is not right for you sex can 'blind' you.
7. Persons and couples with premarital sex experience seem to achieve sexual satisfaction sooner after they are married. However, they are likely to be less satisfied overall with their sex life during marriage. It seems that their premarital sex experiences often rise to haunt them.

Mark Twain once said, "**Sex takes the least amount of time, but causes the most amount of trouble.**" This is profoundly true for teen sex outside of a monogamous marriage. Doctor Dean Ornish proclaims:

I am not aware of any other factor in medicine-not diet, not smoking, not exercise, not stress, not genetics, not drugs, not surgery-that has a greater impact on our quality of life, incidence of illness, and premature death from all causes. Love and intimacy are at the root of what makes us sick and what makes us well, what causes sadness and what brings happiness; what makes us suffer and what leads to healing.

In the 1980s America boasted "safe sex" to alleviate the widespread fear of AIDS. As stated in Joe White's Pure Excitement:



**Condoms fail 100 percent of the time...**  
**in protecting a boy or girl's virginity.**  
**in protecting a girl's reputation.**  
**in protecting a boy's complex sexual memory bank.**  
**in protecting a couple's purity and friendship development.**  
**in protecting a boy's respect for a girl and vice versa.**  
**in protecting a girl or boy's delicate self-image.**

The effects of premarital sexual relations on teenagers protrude into all parts of their lives: mental, emotional, and physical. Self-disciplined youth protect their marriage beds from virus, fungi, bacteria, yeast, spores, and parasitic contamination, and they guard themselves from anger, doubt, stress, brokenness, and heartache. **By choosing to abstain, teenagers free themselves to live life to the fullest without all the burdens of sex outside of marriage.**

# A Parent's Guide to Teen Parties

As a parent, you know the importance of your teen's social life and that parties are a way to socialize and relax. But an unsupervised or poorly planned party can result in unwanted or even tragic consequences. However, parental responsibility is the key to a fun and safe party.

The following is important information from the American Academy of Pediatrics about teen parties.

## Facts about teen parties

- **Guest list.** When a teen plans a party, news spreads very quickly via social networking sites like Facebook or Twitter. Because of these new media, teen parties can grow too large for parents to control.
- **Time and place.** Teen parties often start late at night and move from house to house.

## Facts about alcohol and drugs

**Teens often expect alcohol and marijuana at parties.** Some parents believe that it is better to allow teens to drink in their home so they can keep them safe. While this idea may be well intentioned, it is simply misguided. Parents cannot keep impaired teens safe.

**Alcohol and other drugs impair judgment.** Teens are more likely to have sex, be involved in a violent incident, or suffer an injury after using drugs or alcohol. All too frequently teens die from violence, unintentional injuries, or overdoses related to alcohol and other drugs.

**Alcohol effects teens differently than adults.** For example, compared with adults, teens are more likely to remain awake, to wander about, or to drive a car while having a much greater degree of mental impairment.

## What parents need to know

**Communication and honesty are important to keep your teen safe.** Teens whose parents talk with them regularly about drugs and alcohol are 42% less likely to use substances than those whose parents don't. Tell your teens that you expect them not to use alcohol or other drugs at parties.

**Parent networking is the best prevention tool to combat underage drinking.** Get to know your teen's friends and their parents. If your teen is planning on going to a party, call the parents to ensure that they will be home and that they will not allow drugs or alcohol. If this is not possible, don't let your teen go.

**Parents are legally responsible for anything that happens to a minor who has been served alcohol or other drugs in their home.** If anyone brings alcohol or other drugs to your home, be prepared to contact their parents. And if someone comes to your home already intoxicated, make sure that they get home safely. Help your teen feel responsible for this as well.

## Parents may be criminally or civilly liable if...

- Alcohol is provided to a minor at a party they have organized.
- Someone's property is damaged.
- Someone is injured.
- Someone leaves and gets into a car accident and/or injures someone else.
- Someone dies.
- Understand the local laws about alcohol and other drugs. Laws about alcohol and drug use vary from state to state, so make sure you know what the laws are in your state.

## If you are hosting a teen party...

- Plan in advance.** Go over party plans with your teen. Encourage your teen to plan non-alcohol-related group activities or games.
- Keep parties small.** Ten to 15 teens for each adult. Make sure at least one adult is present at all times. Ask other parents to come over to help you if you need it.
- Set a guest list.** The party should be for invited guests only. No "crashers" allowed. This will help avoid the "open party" situation.
- Set starting and ending times for the party.** Check local curfew laws to determine an ending time.
- Set party "rules" and your expectations.** Discuss them with your teen before the party. Rules should include the following:
  - o No tobacco, alcohol, or other drugs.
  - o No one can leave the party and then return.
  - o Lights are left on at all times.
  - o Certain rooms of the house are off-limits.
- Have plenty of food and non-alcoholic beverages available.** Also, put your alcohol and any prescription or over-the-counter medicines in a locked cabinet.
- Be there, but not square.** Pick out a spot where you can see what is going on without being in the way. You can also help serve snacks and beverages.

## If your teen is going to a party...

- Know where your teen is going and how long he will be there.** Have the phone number and address of the party. Ask your teen to call you if the location of the party changes. Be sure to let your teen know where you will be during the party.
- Call the parent of the party host** to make sure a parent will be home the entire time and supervising the party. Make sure that tobacco, alcohol, and other drugs will not be allowed.
- Talk with your teen beforehand** about how to handle a situation where alcohol is available at a party.

- Make sure your teen has a way to get to and from the party.** Make it easy for your teen to leave a party by making it clear that he can call at any time for a ride home. Discuss why he might need to make such a call. Remind your teen NEVER to ride home with a driver who has been drinking or using other drugs.
- Be up to greet your teen when he comes home.** This can be a good way to check the time and talk about the evening.
- If your teen is **staying overnight** at a friend's house after the party, verify this arrangement with the friend's parents and that they will be home.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

The AAP Committee on Adolescence and AAP Section on Adolescent Health would like to thank the AAP Committee on Substance Abuse and AAP NY Chapter 2 Committee on Adolescence for their assistance in developing this publication.

### From your doctor

The Perkins Pediatric Clinic, LLC  
Michael W. Perkins, MD  
614 North Pine Street  
DeRidder, LA 70634

phone: (337) 462-6000  
[www.perkinspediatrics.com](http://www.perkinspediatrics.com)

